



# **STIC Search Report**

## **Biotech-Chem Library**

**STIC Database Tracking Number: 107731**

**TO: Michael Meller**  
**Location: cm1/10A03**  
**Art Unit: 1654**  
**Wednesday, November 05, 2003**

**Case Serial Number: 10/026408**

**From: Mary Hale**  
**Location: Biotech/Chem Library**  
**CM1-1E01**  
**Phone: 308-4258**

**Mary.Hale@uspto.gov**

### **Search Notes**

**Exmr. Meller-**

Example 1 was searched. Refer to structure drawing, you'll see what components were drawn.

25 structures were retrieved all having one EP patent reference.

Displayed all structures.

Hope this works for you.

Mary

*Mary Hale -- Supervisor, Info. Branch  
STIC - Biotech/Chem. Library  
CM-1 Room E01  
703-308-4258*

# \* Rush Request - Amended <sup>Accession # 107731</sup>

## SEARCH REQUEST FORM

Scientific and Technical Information Center

Requester's Full Name: Michael Miller Examiner #: 69404 Date: 11/5/03  
 Art Unit: 1654 Phone Number 308-4230 Serial Number: 101026, 408  
 Mail Box and Bldg/Room Location: CMI 10 A03 Results Format Preferred (circle): PAPER DISK E-MAIL

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*

Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: \_\_\_\_\_

Inventors (please provide full names): Lerchen, Hans-Georg; Baumgarten, Jörg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas.

Earliest Priority Filing Date: 12/27/2000

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.


please search this specific compound.  
 Do not expand its structure.  
Only this compound highlighted.  
 Example 2: -> see Attached sheet.

1429  
1419

### STAFF USE ONLY

	Type of Search	Vendors and cost where applicable
Searcher: <u>Mary</u>	NA Sequence (#) _____	STN <u>256.11</u>
Searcher Phone #: <u>84858</u>	AA Sequence (#) _____	Dialog _____
Searcher Location: <u>1E11</u>	Structure (#) <u>1</u>	Questel/Orbit _____
Date Searcher Picked Up: _____	Bibliographic _____	Dr Link _____
Date Completed: <u>11/5</u>	Litigation _____	Lexis/Nexis _____
Searcher Prep & Review Time: <u>7</u>	Fulltext _____	Sequence Systems _____
Clerical Prep Time: <u>4</u>	Patent Family _____	WWW/Internet _____
Online Time: <u>10</u>	Other _____	Other (specify) _____

**Example 1:**



5      Educts: II.1, III.1      Procedure: A  
 Yield: 71%       $R_f = 0.28^{(6)}$       [ESI-MS:  $m/e = 1561 = (M+H)^+$ ]

**Example 2: Diastereoisomer A of compound from Example 1**

10      Educts: II.1, III.2      Procedure: A  
 Yield: 32%       $R_f = 0.28^{(6)}$       [ESI-MS:  $m/e = 1561 = (M+H)^+$ ]

Miller  
10/026408

=> dis his


(FILE 'HOME' ENTERED AT 14:19:44 ON 05 NOV 2003)

FILE 'REGISTRY' ENTERED AT 14:19:50 ON 05 NOV 2003

L1 STR  
L2 0 S L1  
L3 25 S L1 FUL

=> d l3 que stat;d 1-25 ide cbib abs

L1 STR



NODE ATTRIBUTES:

NSPEC IS R AT 37  
DEFAULT MLEVEL IS ATOM  
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
NUMBER OF NODES IS 61

STEREO ATTRIBUTES: NONE

L3 25 SEA FILE=REGISTRY SSS FUL L1

100.0% PROCESSED 46 ITERATIONS  
SEARCH TIME: 00.00.01

25 ANSWERS

L3 ANSWER 1 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439865-01-7 REGISTRY  
CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino) carbonyl] amino] phe  
nyl] sulfonyl] amino] phenyl] ethyl] amino] carbonyl] amino] phenyl] amino] thioxome  
thyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-prolyl-, 6-[(4S)-4-ethyl-


Searched by: Mary Hale 308-4258 CM-1 1E01

3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C76 H92 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR19]_q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 2 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439865-00-6 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino) carbonyl] amino] phenyl] sulfonyl] amino] phenyl] ethyl] amino] carbonyl] amino] phenyl] amino] thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucylglycyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C73 H88 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH2CHPhNHCOCH2NHCO-m-C6H4NH[C(:NH)NHR19]q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C6H4SO2NH-m-C6H4CH(CH2CO2H)NHCONH-p-C6H4NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC50 = 40 nM).




L3 ANSWER 3 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-99-0 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino) carbonyl] amino] phe  
 nyl]sulfonyl] amino] phenyl] ethyl] amino] carbonyl] amino] phenyl] amino] thioxome  
 thyl]-L-prolyl-L-leucylglycyl-L-leucyl-L- $\alpha$ -glutamyl-,  
 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C76 H92 N14 O18 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 4 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-98-9 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl)sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-isoleucyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C77 H96 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\text{v}\beta 3$  integrin receptor, e.g., a radical of formula  $\text{R18COCH}_2\text{CHPhNHCOCH}_2\text{NHCO-m-C}_6\text{H}_4\text{NH[C(:NH)NHR19]q}$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\text{v}\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\text{v}\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).


L3 ANSWER 5 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-97-8 REGISTRY

CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[3-[(propylamino) carbonyl] amino] phe  
 nyl]sulfonyl] amino] phenyl] ethyl] amino] carbonyl] amino] phenyl] amino] thioxome  
 thyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-seryl-, 6-[(4S)-4-ethyl-3,4,12,14-  
 tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl]  
 ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C74 H90 N14 O17 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 6 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-96-7 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxome  
 thyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-phenylalanyl-,  
 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C80 H94 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta$ 3 integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta$ 3 integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta$ 3 integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 7 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-95-6 REGISTRY




CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-valyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl]ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C76 H94 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR19]q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).


L3 ANSWER 8 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-94-5 REGISTRY

CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-D-alanyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C74 H90 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\text{v}\beta 3$  integrin receptor, e.g., a radical of formula  $\text{R18COCH}_2\text{CHPhNHCOCH}_2\text{NHCO-m-C}_6\text{H}_4\text{NH}[\text{C}(\text{:NH})\text{NHR19}]_q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\text{v}\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\text{v}\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 9 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-93-4 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-alanyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C74 H90 N14 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 10 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-91-2 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl)sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-prolyl-L-leucylglycyl-L-leucyl-L-asparaginyl-,  
 7-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C80 H98 N16 O18 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).




L3 ANSWER 11 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-90-1 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl)sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-histidyl-, 5-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl]ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C71 H81 N15 O15 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on


human large intestine cell line HT29 (IC50 = 40 nM).

```
L3 ANSWER 12 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN
RN 439864-89-8 REGISTRY
CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-histidylglycyl-L-leucyl-L-asparaginy-,
6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-
pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX
NAME)
FS PROTEIN SEQUENCE; STEREOSEARCH
MF C75 H87 N17 O17 S2
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL
```


\*\*\*RELATED SEQUENCES AVAILABLE WITH SEOLINK\*\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR19]q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on


human large intestine cell line HT29 (IC50 = 40 nM).

L3 ANSWER 13 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-88-7 REGISTRY  
CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolylglycylglycyl-L-leucyl-L-asparaginyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
FS PROTEIN SEQUENCE; STEREOSEARCH  
MF C71 H83 N15 O17 S2  
SR CA  
LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).


L3 ANSWER 14 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-86-5 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-norvalyl-L-asparaginyl-,  
 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C74 H89 N15 O17 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).




L3 ANSWER 15 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-85-4 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-lysyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl]ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C77 H97 N15 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 16 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-84-3 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino) carbonyl] amino] phe  
 nyl]sulfonyl] amino] phenyl] ethyl] amino] carbonyl] amino] phenyl] amino] thioxome  
 thyl]-L-prolyl-L-leucylglycyl-S-methyl-L-cysteinyl-L-histidyl-,  
 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C75 H88 N16 O16 S3  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH2CHPhNHCOCH2NHCO-m-C6H4NH[C(:NH)NHR19]q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C6H4SO2NH-m-C6H4CH(CH2CO2H)NHCONH-p-C6H4NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC50 = 40 nM).

L3 ANSWER 17 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-79-6 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-tryptophyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C82 H95 N15 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\text{v}\beta 3$  integrin receptor, e.g., a radical of formula  $\text{R18COCH}_2\text{CHPhNHCOCH}_2\text{NHCO-m-C}_6\text{H}_4\text{NH}[\text{C}(\text{:NH})\text{NHR19}]_q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\text{v}\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\text{v}\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 18 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-78-5 REGISTRY

CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-S-methyl-L-cysteinyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)

FS PROTEIN SEQUENCE; STEREOSEARCH

MF C75 H92 N14 O16 S3


SR CA

LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta 3$  integrin receptor, e.g., a radical of formula  $R18COCH2CHPhNHCOCH2NHCO-m-C6H4NH[C(:NH)NHR19]q$ , where R18 is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocycloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1; R19 is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C6H4SO2NH-m-C6H4CH(CH2CO2H)NHCONH-p-C6H4NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 ( $IC_{50} = 40 \text{ nM}$ ).

L3 ANSWER 19 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
RN 439864-76-3 REGISTRY




CN L-Valine, 1-[[[4-[[[(1S)-2-carboxy-1-[3-[[[3-  
 [[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]car  
 bonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-  
 asparaginy]-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C75 H91 N15 O17 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocycloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 20 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-75-2 REGISTRY  
 CN L-Valine, 1-[[[4-[[[(1R)-2-carboxy-1-[3-[[[3-  
 [(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]car  
 bonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-  
 asparaginyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C75 H91 N15 O17 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 21 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-74-1 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-asparaginyl-,  
 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C75 H91 N15 O17 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 22 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-73-0 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-, 5-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl]ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C71 H85 N13 O15 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha\beta$ 3 integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha\beta$ 3 integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha\beta$ 3 integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-(PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).




L3 ANSWER 23 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-68-3 REGISTRY  
 CN L-Valine, 1-[[[4-[[[(1S)-2-carboxy-1-[3-[[[3-  
 [[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]car  
 bonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-  
 histidyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C77 H92 N16 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.


AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 24 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-67-2 REGISTRY  
 CN L-Valine, 1-[[[4-[[[(1R)-2-carboxy-1-[3-[[[3-  
 [[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]car  
 bonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-  
 histidyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-  
 pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX  
 NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C77 H92 N16 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


- REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.
- AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v \beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v \beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v \beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

L3 ANSWER 25 OF 25 REGISTRY COPYRIGHT 2003 ACS on STN  
 RN 439864-66-1 REGISTRY  
 CN L-Valine, 1-[[[4-[[[2-carboxy-1-[3-[[[3-[(propylamino)carbonyl]amino]phenyl]sulfonyl]amino]phenyl]ethyl]amino]carbonyl]amino]phenyl]amino]thioxomethyl]-L-prolyl-L-leucylglycyl-L-leucyl-L-histidyl-, 6-[(4S)-4-ethyl-3,4,12,14-tetrahydro-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-4-yl] ester (9CI) (CA INDEX NAME)  
 FS PROTEIN SEQUENCE; STEREOSEARCH  
 MF C77 H92 N16 O16 S2  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXCENTER, USPATFULL


\*\*RELATED SEQUENCES AVAILABLE WITH SEQLINK\*\*


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1: 137:63479 Preparation of conjugates of integrin receptor antagonists and a cytostatic agent having specifically cleavable linking units. Lerchen, Hans-Georg; Baumgarten, Joerg; Lockhoff, Oswald; Albers, Markus; Schoop, Andreas (Bayer Aktiengesellschaft, Germany). Eur. Pat. Appl. EP 1219305 A1 20020703, 127 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR. (English). CODEN: EPXXDW. APPLICATION: EP 2000-128401 20001227.

AB The invention relates to cytostatics CT-LI-Sp-IA [CT denotes a cytotoxic radical or a radical of a cytostatic or a cytostatic derivative which can addnl. carry a hydroxy, carboxy or amino group; LI is a linker group comprising 5- to 8-amino acid residues in the D- or L-configuration, which can each optionally carry protective groups; Sp is absent or a carbonyl or thiocarbonyl radical; IA is a non-peptide radical addressing an  $\alpha v\beta 3$  integrin receptor, e.g., a radical of formula  $R_{18}COCH_2CHPhNHCOCH_2NHCO-m-C_6H_4NH[C(:NH)NHR_{19}]_q$ , where  $R_{18}$  is OH, (un)substituted (cyclo)alkoxy, aryloxy, heterocyclyloxy, a direct bond, or an atom from the group N, O and S, via which the radical is bonded to the rest of the conjugate; q is 0 or 1;  $R_{19}$  is H, (un)substituted (cyclo)alkyl, aryl, heterocyclyl, an alkylamine or alkylamide radical, or a direct bond, via which the radical is bonded to the rest of the conjugate] and their physiol. acceptable salts and stereoisomers. The cytostatics have a tumor-specific action as a result of linkage to  $\alpha v\beta 3$  integrin antagonists via preferred linking units which can be selectively cleaved by enzymes such as metallo matrix proteases (MMPs5), i.e., by enzymes which can especially be found in tumor tissue. The preferred linking units guarantee the serum stability of the conjugate of cytostatic and  $\alpha v\beta 3$  integrin antagonist and, at the same time, the desired intracellular action within tumor cells as a result of its specific enzymic or hydrolytic cleavability with release of the cytostatic. Thus, 20-O-[PrNHCONH-m-C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NH-m-C<sub>6</sub>H<sub>4</sub>CH(CH<sub>2</sub>CO<sub>2</sub>H)NHCONH-p-C<sub>6</sub>H<sub>4</sub>NHC(S)-Pro-Leu-Gly-Leu-His-Val]camptothecin (1) was prepared by reaction of 20(S)-camptothecin with N-(tert-butoxycarbonyl)-L-valine-N-carboxyanhydride, deprotection, peptide coupling reactions, and formation of the thiourea linkage. Compound 1 was assayed for cytostatic action on human large intestine cell line HT29 (IC<sub>50</sub> = 40 nM).

=> fil caol;s l3  
COST IN U.S. DOLLARS

	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	252.85	253.06

	SINCE FILE	TOTAL
	ENTRY	SESSION
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)		
CA SUBSCRIBER PRICE	-15.50	-15.50

FILE 'CAOLD' ENTERED AT 14:28:27 ON 05 NOV 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1907-1966  
FILE LAST UPDATED: 01 May 1997 (19970501/UP)

This file contains CAS Registry Numbers for easy and accurate substance identification. Title keywords, authors, patent assignees, and patent information, e.g., patent numbers, are now searchable from 1907-1966. TIFF images of CA abstracts printed between 1907-1966 are available in the PAGE display formats.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

L4 0 L3

=> fil hcaplus;s l3  
COST IN U.S. DOLLARS

	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.40	253.46

	SINCE FILE	TOTAL
	ENTRY	SESSION
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)		
CA SUBSCRIBER PRICE	0.00	-15.50

FILE 'HCAPLUS' ENTERED AT 14:28:45 ON 05 NOV 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 5 Nov 2003 VOL 139 ISS 19  
FILE LAST UPDATED: 4 Nov 2003 (20031104/ED)

Searched by: Mary Hale 308-4258 CM-1 1E01

This file contains CAS Registry Numbers for easy and accurate substance identification.

L5                    1 L3

=> fil reg;save l1 meller/q

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	2.25	255.71
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-15.50

FILE 'REGISTRY' ENTERED AT 14:29:07 ON 05 NOV 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES:     3 NOV 2003    HIGHEST RN 612478-18-9  
DICTIONARY FILE UPDATES:    3 NOV 2003    HIGHEST RN 612478-18-9

TSCA INFORMATION NOW CURRENT THROUGH JULY 14, 2003

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

QUERY L1 HAS BEEN SAVED AS 'MELLER/Q'

=> log y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.40	256.11
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-15.50

STN INTERNATIONAL LOGOFF AT 14:29:10 ON 05 NOV 2003